

# WESTSIDE GASTROENTEROLOGY

## CONSULTANTS

### **Colonoscopy Bowel Preparation Instructions (SUPREP®)**

Note that these instructions are also used if you are having an upper endoscopy and a colonoscopy scheduled for the same day.

#### **PLEASE READ ALL INSTRUCTIONS ON THE DAY YOU RECEIVE THEM**

Our goal is to provide excellent quality care during your visit with us and throughout your experience with our office. We have included some very important information pertaining to your procedure and care.

#### **About Colonoscopy**

Bowel preparation (cleansing) is needed to perform effective colonoscopy. Any stool remaining in the colon can hide lesions and result in the need to repeat the colonoscopy. You should plan to be at the procedure area 3-4 hours. It is critical that you follow the instructions as directed. Your physician will discuss your procedure results with you and your family members when you are in the recovery room. If any biopsies or tissue removal is performed, you will receive a letter with those results about 2-3 weeks after the procedure. If there are serious findings on the biopsy, your physician will contact you.

Every effort will be made to keep your appointment at the schedule time, but unexpected delays and emergencies may occur and affect your procedure time. We give each patient the attention needed for his or her procedure and apologize in advance for delays if these occur.

#### **14 DAYS BEFORE PROCEDURE**

- If you are taking weight loss medication, contact your prescribing physician for instructions on stopping the medication prior to your procedure.

## **7 DAYS BEFORE PROCEDURE**

- Read all prep instructions.
- Purchase prep medications and clear liquid diet items.
- **Make arrangements for someone 18 years or older to come with you to the procedure.** Sedation is usually given during your procedure. A responsible adult must be present before, during and after your procedure. Drop offs are not allowed. After your procedure you will not be able to drive, operate machinery, make important decisions, or return to work for the rest of the day. You may resume normal activities the next day unless your doctor directs otherwise. No Taxi or Uber/Lyft drivers.
- **Begin avoiding corn, popcorn, nuts, or foods containing visible seeds.**
- Stop any multi-vitamins, iron supplements, and herbal supplements.

## **2-7 DAYS BEFORE PROCEDURE**

- **STOP anticoagulants/blood thinners** exactly as you have been instructed by your prescribing physician. This may include clopidogrel (Plavix<sup>®</sup>), warfarin (Coumadin<sup>®</sup>, Jantoven<sup>®</sup>), dabigatran (Pradaxa<sup>®</sup>), apixaban (Eliquis<sup>®</sup>) or other non-aspirin blood thinners. Do not stop these medications on your own. If you are uncertain if you need to stop a medication, please call our office for instructions. If you take low dose aspirin (81 mg), you may continue taking it.
- If you have diabetes and were not given specific instructions for management of medications, call our office for instructions.

## **2 DAYS BEFORE PROCEDURE**

- Stop anti-inflammatory medications ibuprofen (Motrin<sup>®</sup>, Advil<sup>®</sup>), naproxen (Aleve<sup>®</sup>).
- Celebrex<sup>®</sup> and acetaminophen (Tylenol<sup>®</sup>) are ok to use.

## **1 DAY BEFORE PROCEDURE**

- No solid food.
- Drink plenty of fluids throughout the day to avoid dehydration including: water, clear fruit juices, white grape juice, apple juice, Kool-Aid®, POWERADE®, clear soup broth or bouillon, popsicles, coffee (no cream), tea, hard candy/jelly beans, soda pop (7-Up®, Sprite®, regular or diet Pepsi® and Coke®, ginger ale, orange soda), Jell-O® (without fruit or other additions).
- **NO ALCOHOL**
- **NO RED/PURPLE colored drinks, popsicles, or Jell-O®.**
- At 6pm, take your **first dose of SUPREP®**. Use the provided cup to mix with Gatorade®, Ginger Ale or Sprite®. Drink it cold and slow. One hour later, drink 2 cups of water.

## **DAY OF PROCEDURE**

- Take your **second dose of SUPREP® 4-6 hours prior to arrival time.**
- Take it the same way as the first dose by mixing with Gatorade®, Ginger Ale, or Sprite®.
- Do not eat or drink anything (SOLID, LIQUID OR ALCOHOL) except the prep solution 4-6 hours prior to arrival time.
- You can take your morning medications with sips of water but do not take ace inhibitor blood pressure medication (e.g. lisinopril, enalapril, benazepril).
- Wear loose fitting clothes, NO jewelry, makeup, or hair products

The goal of the bowel prep is to have colorless (or close to colorless) liquid stool by the time the colonoscopy procedure will occur. It is normal to have yellow or green diarrhea, regardless of the color of the clear liquid diet drinks.

**Note:** If you think the prep is not working call the office at 317-745-7310, or if after hours call the hospital operator at 317-745-4451 and ask for the GI doctor on call.

## **What to expect after the procedure**

- Check your discharge paperwork for instructions regarding new medications or changes to your prior medications and other directions.
- You may also experience some cramping or bloating because of the air introduced into the colon during the procedure. This should disappear quickly when you pass gas. You may consider taking Gas-X® pills (over the counter) at home to help with this. Call our office with any severe or worsening abdominal pain.
- You were given sedation during the procedure. A responsible adult must drive you home. No driving is allowed for the rest of the day. No exceptions.
- We recommend having a responsible adult stay with you through the rest of the day after the procedure.
- Rest at home with activity as tolerated. Avoid dangerous activities.
- Even if you feel alert after the procedure your judgment and coordination may be impaired for the rest of the day. No alcohol, as this can interact with sedation you received.
- After your procedure, progress as tolerated to your usual diet. You may want to start with a bland diet such as bread, rice, pudding, applesauce, mashed potatoes, cold cereal, or ice cream.
- The day following your procedure you may resume all normal activities and diet.

**If you must cancel or reschedule your procedure, please call Endoscopy Scheduling at 317-745-7310 as soon as possible. If you do not cancel your appointment with sufficient notice, you may be dismissed from your office/practice.**

**Please contact Westside Gastroenterology at 317-745-7310 with any questions or concerns.**

Here are some educational resources that you may find helpful:

<https://www.youtube.com/watch?v=eA1PIMa1ULg>

<https://gastro.org/practice-guidance/gi-patient-center/topic/colonoscopy/>

American College of Gastroenterology <https://gi.org/>

American Society of Gastrointestinal Endoscopy <https://www.asge.org/>

American Gastroenterological Association <https://gastro.org/>