WESTSIDE GASTROENTEROLOGY CONSULTANTS

Colonoscopy Bowel Preparation Instructions for 2 Days Prior to Procedure (14 M) MiraLAX[®]

PLEASE READ ALL INSTRUCTIONS ON THE DAY YOU RECEIVE THEM

Bowel Prep and Clear Liquids Shopping List

These medications are all available over the counter without a prescription.

- MiraLax[®] 238-gram bottle (Enough for 14 doses). Other names are Glycolax[®] or Polyethylene Glycol 3350 - these are identical to MiraLax[®].
- Bisacodyl laxative tablets (common brand name is Dulcolax[®]) 4 tablets each day (8) will be needed.
- TWO (2) 28 oz bottles of Gatorade[®] (NO RED or PURPLE drinks). Patients who do not like Gatorade[®] or have diabetes should use G2[®] (low sugar), Gatorade Zero[®], POWERADE Zero[®] or Pedialyte[®].
- <u>OPTIONAL</u> **Simethicone** 80mg or 125mg (common brand name is Gas-X[®]) to use if needed for bloating symptoms.
- <u>OPTIONAL</u> a **petroleum-based ointment product** or diaper rash ointment for potential irritation from frequent bowel movements.

2 DAYS BEFORE PROCEDURE

- No solid food
- May have full liquids (e. g. ice cream, strained cream soups, milk, yogurt, sherbet, milkshakes, custard, pudding)
- Drink plenty of fluids throughout the day to avoid dehydration including water, clear fruit juices, white grape juice, apple juice, Kool-Aid[®], POWERADE[®], clear soup broth or bouillon, popsicles, coffee (no cream), tea, hard candy/jelly beans, soda pop (7-Up[®], Sprite[®], regular or diet Pepsi[®] and Coke[®], ginger ale, orange soda), Jell-O[®] (without fruit or other additions).
- NO ALCOHOL
- NO RED/PURPLE colored drinks, popsicles, or Jell-O®

- STEP ONE: Mix 7 capfuls of Miralax[®] into each 28-ounce Gatorade[®] bottle until dissolved.
- STEP TWO: At approximately 4 pm, take 4 Dulcolax[®] laxative tablets with water.
- STEP THREE: At approximately 6 pm begin drinking the first bottle of the Miralax[®] /Gatorade[®] solution.
- STEP FOUR: Drink the second Miralax[®]/Gatorade[®] bottle before bedtime.